

# Wound signaling: The missing link in plant regeneration

Lyuqin Chen, Beibei Sun, Lin Xu & Wu Liu

To cite this article: Lyuqin Chen, Beibei Sun, Lin Xu & Wu Liu (2016) Wound signaling: The missing link in plant regeneration, Plant Signaling & Behavior, 11:10, e1238548, DOI: [10.1080/15592324.2016.1238548](https://doi.org/10.1080/15592324.2016.1238548)

To link to this article: <https://doi.org/10.1080/15592324.2016.1238548>



Published online: 23 Sep 2016.



Submit your article to this journal [↗](#)



Article views: 1970



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 12 View citing articles [↗](#)

ARTICLE ADDENDUM

## Wound signaling: The missing link in plant regeneration

Lyuqin Chen<sup>a,b</sup>, Beibei Sun<sup>a,c</sup>, Lin Xu<sup>a</sup>, and Wu Liu<sup>a</sup>

<sup>a</sup>National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, China; <sup>b</sup>University of Chinese Academy of Sciences, Beijing, China; <sup>c</sup>College of Life and Environment Sciences, Shanghai Normal University, Shanghai, China

### ABSTRACT

Wounding is the first event that occurs in plant regeneration. However, wound signaling in plant regeneration is barely understood. Using a simple system of *de novo* root organogenesis from *Arabidopsis thaliana* leaf explants, we analyzed the genes downstream of wound signaling. Leaf explants may produce at least two kinds of wound signals to trigger short-term and long-term wound signaling. Short-term wound signaling is primarily involved in controlling auxin behavior and the fate transition of regeneration-competent cells, while long-term wound signaling mainly modulates the cellular environment at the wound site and maintains the auxin level in regeneration-competent cells. *YUCCA* (*YUC*) genes, which are involved in auxin biogenesis, are targets of short-term wound signaling in mesophyll cells and of long-term wound signaling in regeneration-competent cells. The expression patterns of *YUCs* provide important information about the molecular basis of wound signaling in plant regeneration.

### ARTICLE HISTORY

Received 13 September 2016  
Accepted 15 September 2016

### KEYWORDS

*Arabidopsis thaliana*; *De novo* root organogenesis; plant regeneration; wound signals; wound signaling


Plant cells are highly plastic and have powerful regenerative abilities.<sup>1–4</sup> Wounding is the first event in plant regeneration.<sup>1, 5</sup> Studies on wounding have suggested several candidates for wound signals, such as electrical pulses, hydraulic pressure, Ca<sup>2+</sup>, reactive oxygen species, the oligopeptide systemin, oligosaccharides, jasmonic acid, salicylic acid, ethylene, abscisic acid, and changes in various metabolic processes.<sup>6,7</sup> Many studies have tried to clarify the effects of wounding on regeneration by analysis of the genes downstream of wound signaling.<sup>8–27</sup> However, our knowledge about whether and how the above physical and chemical signals serve as wound signal (s) in plant regeneration is still limited.

To study the role of wound signaling in *de novo* root organogenesis, we cultured leaf explants on B5 medium to regenerate adventitious roots.<sup>28</sup> Wounding has complex biological effects, and may have multiple roles in *de novo* root organogenesis from leaf explants. Our previous studies revealed that at least two pathways are triggered by wounding of leaf explants: short-term and long-term wound-signaling pathways.

Short-term wound signaling is required for auxin production and cell fate transition (indicated by the red flow path in Fig. 1A).<sup>10,29</sup> This pathway comprises at least three stages of signal delivery (Fig. 1A). In stage I, wounding first triggers short-term wound signaling that lasts from seconds to hours. This wound signal spreads very rapidly from the wound site to mesophyll cells and activates *YUCCA1* (*YUC1*) and *YUC4* expression in mesophyll cells within 4 hours. Stage II involves auxin behavior. Auxin begins to be produced by *YUCs* in mesophyll cells within 4 hours and then polar-transported to regeneration-competent cells in the vasculature near the wound site at around 12 hours after wounding. In stage III, the

accumulation of auxin in regeneration-competent cells activates expression of *WUSCHEL RELATED HOMEODOMAIN 11* (*WOX11*) and *WOX12*, which start the cell fate transition to form root founder cells at around 1 to 2 days d after wounding.

The long-term wound-signaling pathway functions for a relatively long time, for around 2 days after wounding (indicated by the blue flow path in Fig. 1A).<sup>11</sup> One event downstream of long-term wound signaling is the activation of a group of *NAC* (*NAM*, *ATAF1,2*, and *CUC2*) transcription factor genes including *NAC1* (Fig. 1A).<sup>11</sup> *NAC1* is expressed in many cells at the wound site around 1 to 2 days after detachment of the leaf explant. However, *NAC1* does not affect the fate transition of regeneration-competent cells because *WOX11* and *WOX12* expression are normally activated when *NAC1* function is blocked. The role of *NAC1* might be to control the cellular environment, including cell wall metabolism. On the other hand, the long-term wound-signaling pathway also activates *YUC4* expression in regeneration-competent cells near the wound site at around 2 days after wounding (Fig. 1A),<sup>10</sup> and this might contribute to maintaining a high auxin level in regeneration-competent cells. Here, we analyzed *YUC4<sub>pro</sub>:GUS*<sup>10</sup> in leaf explants cultured on B5 medium containing naphthylphthalamic acid (NPA, a polar auxin transport inhibitor) to test whether *YUC4* is a direct target of long-term wound signaling or is activated by auxin accumulation in regeneration-competent cells. Treatment with NPA can prevent auxin accumulation in regeneration-competent cells in leaf explants.<sup>29</sup> We observed strong  $\beta$ -glucuronidase (*GUS*) signals in regeneration-competent cells near the wound in the leaf explant cultured on medium containing NPA at 2 days after culture



**Figure 1.** Wound signaling in *de novo* root organogenesis. (A) Model of short-term (red flow path) and long-term (blue flow path) wound signaling-mediated pathways in *de novo* root organogenesis. Orange region indicates wound site. (B) *YUC4* expression in leaf explant cultured on B5 medium containing 1  $\mu$ M NPA. Arrow shows GUS signal in regeneration-competent cells at 2 DAC. Leaf explant was cultured in dark conditions,<sup>28</sup> and GUS staining was performed as described previously.<sup>29,30</sup> Scale bar, 500  $\mu$ m in (B).

(DAC) (Fig. 1B), suggesting that *YUC4* might be a direct target of long-term wound signaling in regeneration-competent cells. Currently, it is unclear whether the activation of *NAC1* and *YUC4* is controlled dependently or independently by long-term wound signaling.

The short-term and long-term wound-signaling pathways act differently in leaf explants. First, the short-term wound signal is produced at the wound site and quickly moves to the mesophyll cells to activate *YUC1* and *YUC4* expression.<sup>10</sup> Therefore, this signal might be a moving signal that can spread from cell to cell. Long-term wound signaling functions at the wound site,<sup>11</sup> suggesting that it is a local signal. Second, short-term wound signaling lasts for seconds to hours,<sup>10</sup> while long-term signaling lasts for at least 2 days.<sup>11</sup> This suggests that the short-term wound signal has a short lifetime, while the long-term signal has a longer lifetime or is continuously produced in response to wounding. Third, short-term wound signaling seems to function in both the detached explant and in the wounded leaf residue on the source plant, because *YUC1* and *YUC4* expression are quickly activated in leaf explants as well as at the wound site on the source plant.<sup>10</sup> However, long-term wound signaling activates *NAC1* expression primarily at the wound site on the detached explant, and not in the leaf residue on the source plant.<sup>11</sup> This indicates that long-term wound signaling has a role in distinguishing different types of wounding. Overall, it seems that wounding produces complex molecules that serve as different types of wound signals in *de novo* root organogenesis from leaf explants.

Although many of the downstream effects of wounding have been identified, our knowledge about wound signals and wound signaling at the molecular level is still very limited. Recent studies have suggested that jasmonic acid, ethylene, and

electrical pulses have effects on tissue repair,<sup>9,15,17,25</sup> raising the possibility that they may be candidates for wound signals to trigger regeneration. In our regeneration system of *de novo* root organogenesis, *YUC1* and *YUC4* serve as targets of short-term wound signaling in mesophyll cells, and *NAC1* and *YUC4* are targets of long-term wound signaling at the wound site. It will be interesting to test the upstream molecular events of *YUCs* and *NAC1* to study the molecular basis of wounding during *de novo* root organogenesis.<sup>18</sup>

## Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

## Funding

This work was supported by grants from National Basic Research Program of China (973 Program, 2014CB943500/2012CB910503), the National Natural Science Foundation of China (91419302/31422005) and Youth Innovation Promotion Association CAS (2014241).

## References

- Xu L, Huang H. Genetic and epigenetic controls of plant regeneration. *Curr Top Dev Biol* 2014; 108:1-33; PMID:24512704; <http://dx.doi.org/10.1016/B978-0-12-391498-9.00009-7>
- Ikeuchi M, Ogawa Y, Iwase A, Sugimoto K. Plant regeneration: cellular origins and molecular mechanisms. *Development* 2016; 143:1442-51; PMID:27143753; <http://dx.doi.org/10.1242/dev.134668>
- Pulianmackal AJ, Kareem AV, Durgaprasad K, Trivedi ZB, Prasad K. Competence and regulatory interactions during regeneration in plants. *Front Plant Sci* 2014; 5:142; PMID:24782880; <http://dx.doi.org/10.3389/fpls.2014.00142>

4. Su YH, Zhang XS. The hormonal control of regeneration in plants. *Curr Top Dev Biol* 2014; 108:35-69; PMID:24512705; <http://dx.doi.org/10.1016/B978-0-12-391498-9.00010-3>
5. Lup SD, Tian X, Xu J, Perez-Perez JM. Wound signaling of regenerative cell reprogramming. *Plant Sci* 2016; 250:178-87; PMID:27457994; <http://dx.doi.org/10.1016/j.plantsci.2016.06.012>
6. Leon J, Rojo E, Sanchez-Serrano JJ. Wound signalling in plants. *J Exp Bot* 2001; 52:1-9; PMID:11181708; <http://dx.doi.org/10.1093/jexbot/52.354.1>
7. Maffei ME, Mithofer A, Boland W. Before gene expression: early events in plant-insect interaction. *Trends Plant Sci* 2007; 12:310-6; PMID:17596996; <http://dx.doi.org/10.1016/j.tplants.2007.06.001>
8. Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M, Sakakibara H, Sugimoto K, et al. The AP2/ERF transcription factor WIND1 controls cell dedifferentiation in *Arabidopsis*. *Curr Biol* 2011; 21:508-14; PMID:21396822; <http://dx.doi.org/http://dx.doi.org/10.1016/j.cub.2011.02.020>
9. Melnyk CW, Schuster C, Leyser O, Meyerowitz EM. A Developmental Framework for Graft Formation and Vascular Reconnection in *Arabidopsis thaliana*. *Curr Biol* 2015; 25:1306-18; PMID:25891401; <http://dx.doi.org/10.1016/j.cub.2015.03.032>
10. Chen L, Tong J, Xiao L, Ruan Y, Liu J, Zeng M, Huang H, Wang JW, Xu L. YUCCA-mediated auxin biogenesis is required for cell fate transition occurring during *de novo* root organogenesis in *Arabidopsis*. *J Exp Bot* 2016; 67:4273-84; PMID:27255928; <http://dx.doi.org/10.1093/jxb/erw213>
11. Chen X, Cheng J, Chen L, Zhang G, Huang H, Zhang Y, Xu L. Auxin-independent NAC pathway acts in response to explant-specific wounding and promotes root tip emergence during *de novo* root organogenesis in *Arabidopsis*. *Plant Physiol* 2016; 170:2136-45; PMID:26850273; <http://dx.doi.org/10.1104/pp.15.01733>
12. Efroni I, Mello A, Navy T, Ip PL, Rahni R, DelRose N, Powers A, Sattija R, Birnbaum KD. Root Regeneration Triggers an Embryo-like Sequence Guided by Hormonal Interactions. *Cell* 2016; 165:1721-33; PMID:27212234; <http://dx.doi.org/10.1016/j.cell.2016.04.046>
13. Xu J, Hofhuis H, Heidstra R, Sauer M, Friml J, Scheres B. A molecular framework for plant regeneration. *Science* 2006; 311:385-8; PMID:16424342; <http://dx.doi.org/10.1126/science.1121790>
14. Sena G, Wang X, Liu HY, Hofhuis H, Birnbaum KD. Organ regeneration does not require a functional stem cell niche in plants. *Nature* 2009; 457:1150-3; PMID:19182776; <http://dx.doi.org/10.1038/nature07597>
15. Asahina M, Azuma K, Pitaksaringkarn W, Yamazaki T, Mitsuda N, Ohme-Takagi M, Yamaguchi S, Kamiya Y, Okada K, Nishimura T, et al. Spatially selective hormonal control of RAP2.6L and ANAC071 transcription factors involved in tissue reunion in *Arabidopsis*. *Proc Natl Acad Sci U S A* 2011; 108:16128-32; PMID:21911380; <http://dx.doi.org/10.1073/pnas.1110443108>
16. Ahkami AH, Lischewski S, Haensch KT, Porfirova S, Hofmann J, Roll-etschek H, Melzer M, Franken P, Hause B, Druege U, et al. Molecular physiology of adventitious root formation in *Petunia hybrida* cuttings: involvement of wound response and primary metabolism. *New Phytol* 2009; 181:613-25; PMID:19076299; <http://dx.doi.org/10.1111/j.1469-8137.2008.02704.x>
17. Kral N, Ougolnikova AH, Sena G. Externally imposed electric field enhances plant root tip regeneration. *Regeneration* 2016; 3(3):156-67; PMID:27606066; <http://dx.doi.org/10.1002/reg2.59>
18. Sang YL, Cheng ZJ, Zhang XS. Endogenous auxin biosynthesis and *de novo* root organogenesis. *J Exp Bot* 2016; 67:4011-3; PMID:27402616; <http://dx.doi.org/10.1093/jxb/erw250>
19. Delessert C, Wilson IW, Van Der Straeten D, Dennis ES, Dolferus R. Spatial and temporal analysis of the local response to wounding in *Arabidopsis* leaves. *Plant Mol Biol* 2004; 55:165-81; PMID:15604673; <http://dx.doi.org/10.1007/s11103-004-0112-7>
20. Sukumar P, Maloney GS, Muday GK. Localized induction of the ATP-binding cassette B19 auxin transporter enhances adventitious root formation in *Arabidopsis*. *Plant Physiol* 2013; 162:1392-405; PMID:23677937; <http://dx.doi.org/10.1104/pp.113.217174>
21. Chupeau MC, Granier F, Pichon O, Renou JP, Gaudin V, Chupeau Y. Characterization of the early events leading to totipotency in an *Arabidopsis* protoplast liquid culture by temporal transcript profiling. *Plant Cell* 2013; 25:2444-63; PMID:23903317; <http://dx.doi.org/10.1105/tpc.113.109538>
22. Pitaksaringkarn W, Ishiguro S, Asahina M, Satoh S. ARF6 and ARF8 contribute to tissue reunion in incised *Arabidopsis* inflorescence stems. *Plant Biotechnol J* 2014; 34:49-53; <http://dx.doi.org/10.5511/plantbiotechnology.13.1028b>
23. Asahina M, Satoh S. Molecular and physiological mechanisms regulating tissue reunion in incised plant tissues. *J Plant Res* 2015; 128:381-8; PMID:25736731; <http://dx.doi.org/10.1007/s10265-015-0705-z>
24. Grafi G, Barak S. Stress induces cell dedifferentiation in plants. *Biochim Biophys Acta* 2015; 1849:378-84; PMID:25086338; <http://dx.doi.org/10.1016/j.bbagr.2014.07.015>
25. Reid JB, Ross JJ. Regulation of tissue repair in plants. *Proc Natl Acad Sci U S A* 2011; 108:17241-2; PMID:21960442; <http://dx.doi.org/10.1073/pnas.1114432108>
26. Heyman J, Cools T, Vandenbussche F, Heyndrickx KS, Van Leene J, Vercauteren I, Vanderauwera S, Vandepoele K, De Jaeger G, Van Der Straeten D, et al. ERF115 controls root quiescent center cell division and stem cell replenishment. *Science* 2013; 342:860-3; PMID:24158907; <http://dx.doi.org/10.1126/science.1240667>
27. Ge Y, Liu J, Zeng M, He J, Qin P, Huang H, Xu L. Identification of WOX family genes in *Selaginella kraussiana* for studies on stem cells and regeneration in lycophytes. *Front Plant Sci* 2016; 7:93; PMID:26904063; <http://dx.doi.org/10.3389/fpls.2016.00093>
28. Chen X, Qu Y, Sheng L, Liu J, Huang H, Xu L. A simple method suitable to study *de novo* root organogenesis. *Front Plant Sci* 2014; 5:208; PMID:24860589; <http://dx.doi.org/10.3389/fpls.2014.00208>
29. Liu J, Sheng L, Xu Y, Li J, Yang Z, Huang H, Xu L. WOX11 and 12 are involved in the first-step cell fate transition during *de novo* root organogenesis in *Arabidopsis*. *Plant Cell* 2014; 26:1081-93; PMID:24642937; <http://dx.doi.org/10.1105/tpc.114.122887>
30. He C, Chen X, Huang H, Xu L. Reprogramming of H3K27me3 is critical for acquisition of pluripotency from cultured *Arabidopsis* tissues. *PLoS Genet* 2012; 8:e1002911; PMID:22927830; <http://dx.doi.org/10.1371/journal.pgen.1002911>